

CLINICAL SCIENCE OF NUTRITION

VOLUME 2
ISSUE 1
APRIL 2020

Editor in Chief

Sadık Kılıçturgay

Department of General Surgery, Uludağ University School of Medicine, Bursa, Turkey

ORCID ID: 0000-0002-2427-8344

Associate Editors

R. Haldun Gündoğdu

Department of Gastrointestinal Surgery, Ankara Şehir Hastanesi, Ankara, Turkey

ORCID ID: 0000-0002-7021-4827

Mehmet Uyar

Department of Anesthesiology and Reanimation, Ege University School of Medicine, İzmir, Turkey

ORCID ID: 0000-0001-9292-2616

Consultant in Biostatistics

Şule Oktay

Kappa Consultancy Training Research, İstanbul Turkey

Advisory Board

Sedat Boyacıoğlu

Department of Gastroenterology, Başkent University School of Medicine, Ankara, Turkey

İsmail Cinel

Department of Anesthesiology and Reanimation, Marmara University School of Medicine, İstanbul, Turkey

Rüksan Çehreli

Department of Preventive Oncology, Institute of Oncology, Dokuz Eylül University School of Medicine, İzmir, Turkey

Seher Demirer

Department of General Surgery, Ankara University School of Medicine, Ankara, Turkey

Meltem Gülhan Halil

Department of Geriatrics, Hacettepe University School of Medicine, Ankara, Turkey

Kürşat Gündoğan

Department of Intensive Care, Erciyes University School of Medicine, Kayseri, Turkey

Levent Güngör

Department of Neurology, Ondokuz Mayıs University School of Medicine, Samsun, Turkey

Diclehan Kılıç

Department of Radiation Oncology, Gazi University School of Medicine, Ankara, Turkey

Gül Kızıltan

Department of Nutrition and Dietetics, Başkent University Faculty of Health Sciences, Ankara, Turkey

Hasan Özen

Department of Pediatrics, Hacettepe University School of Medicine, Ankara, Turkey

Bülent Saka

Department of Internal Diseases, İstanbul University, İstanbul School of Medicine, İstanbul, Turkey

Ferda Şöhret Kahveci

Department of Anesthesiology and Reanimation, Uludağ University School of Medicine, Bursa, Turkey

Tuğba Yavuzşen

Department of Medical Oncology, Dokuz Eylül University School of Medicine, İzmir, Turkey

Kaya Yorgancı

Department of General Surgery, Hacettepe University School of Medicine, Ankara, Turkey



Publisher

İbrahim KARA

Publication Director

Ali ŞAHİN

Editorial Development

Gizem KAYAN TEKAÜT

Deputy Publication Director

Gökhan ÇİMEN

Publication Coordinators

İrem SOYSAL

Arzu YILDIRIM

Deniz KAYA

Gülnur MERCAN

Bahar ALBAYRAK

Project Coordinators

Sinem KOZ

Doğan ORUÇ

Finance and Administration

Zeynep YAKIŞIRER ÜREN

Betül ÇİMEN

Graphics Department

Ünal ÖZER

Deniz Elif DURAN

Beyzanur KARABULUT

Contact:

Address: Büyükdere Cad. 105/9 34394

Mecidiyeköy, Şişli, İstanbul

Phone: +90 212 217 17 00

Fax : +90 212 217 22 92

E-mail: info@avesyayincilik.com

AIMS AND SCOPE

Clinical Science of Nutrition (Clin Sci Nutr) is an international, scientific, open Access periodical published in accordance with independent, unbiased, and double-blinded peer-review principles. The journal is the official publication of the Society of Clinical Enteral Parenteral Nutrition – Turkey, and it is published tri-annually in April, August, and December. The publication language of the journal is English.

The journal aims to contribute to the literature by publishing high impact content and become one of the leading publications of the field while functioning as an open discussion forum on significant issues of current interest. Clinical Science of Nutrition also aims to have significant input in emphasizing the increasing importance of clinical nutrition in Turkey and the region, identifying the effects of differences between societies on study results in a clearer way and converting clinical applications into scientific publications as well as forming a bridge between West and East.

The scope of Clinical Science of Nutrition includes original research articles, review articles, case reports, conference reports, and letters to the editor as well as editorials, abstracts from international and national congresses, panel meetings, conferences and symposia. As an online-only publication, in addition to traditional manuscript submissions, Clinical Science of Nutrition is also able to process video, audio and interactive software submissions. Authors are encouraged to submit their content in the most appropriate medium to best convey their findings to the audience of Clinical Science of Nutrition.

The journal covers all aspects of nutrition and dietetics including prevalence of malnutrition and its effects on clinical results; nutritional support and delivery methods and their advantages and disadvantages; nutritional support products and their side effects; immune system and nutritional support; ERAS protocol and nutritional support; home parenteral and enteral nutrition; nutrition support teams and their necessity, challenges and potential solutions of nutritional support.

The journal's target audience includes academicians, practitioners, specialists and students interested in nutrition and dietetics.

The editorial and publication processes of the journal are shaped in accordance with the guidelines of the International Committee of Medical Journal Editors (ICMJE), World Association of Medical Editors (WAME), Council of Science Editors (CSE), Committee on Publication Ethics (COPE), European Association of Science Editors (EASE), and National Information Standards Organization (NISO). The journal is in conformity with the Principles of Transparency and Best Practice in Scholarly Publishing (doaj.org/bestpractice).

Processing and publication are free of charge with the journal. No fees are requested from the authors at any point throughout the evaluation and publication process. All manuscripts must be submitted via the online submission system, which is available at clinscinutr.org. The journal guidelines, technical information, and the required forms are available on the journal's web page.

Publication expenses of the journal are covered by the Society of Clinical Enteral Parenteral Nutrition - Turkey. Potential advertisers should contact the Editorial Office. Advertisement images are published only upon the Editor-in-Chief's approval.

Statements or opinions expressed in the manuscripts published in the journal reflect the views of the author(s) and not the opinions of the Society of Clinical Enteral Parenteral Nutrition - Turkey, editors, editorial board, and/or publisher; the editors, editorial board, and the publisher disclaim any responsibility or liability for such materials.

All published content is available online, free of charge at clinscinutr.org.

Clinical Science of Nutrition is an open access publication and the journal's publication model is based on Budapest Open Access Initiative (BOAI) declaration. Journal's archive is available online, free of charge at clinscinutr.org. Clinical Science of Nutrition's content is licensed under a Creative Commons Attribution-Non-Commercial 4.0 International License.



Editor in Chief: Sadık Kılıçturgay

Address: Department of General Surgery, Uludağ University School of Medicine, Bursa, Turkey

E-mail: skturgay@gmail.com

Publisher: AVES

Address: Büyükdere Cad. 105/9 34394 Mecidiyeköy, Şişli, İstanbul, Turkey

Phone: +90 212 217 17 00

Fax: +90 212 217 22 92

E-mail: info@avesyayincilik.com

Web page: avesyayincilik.com

INSTRUCTIONS TO AUTHORS

Clinical Science of Nutrition (Clin Sci Nutr) is an international, scientific, open Access periodical published in accordance with independent, unbiased, and double-blinded peer-review principles. The journal is the official publication of the Society of Clinical Enteral Parenteral Nutrition – Turkey, and it is published tri-annually in April, August, and December. The publication language of the journal is English.

The journal aims to contribute to the literature by publishing high impact content and become one of the leading publications of the field while functioning as an open discussion forum on significant issues of current interest. Clinical Science of Nutrition also aims to have significant input in emphasizing the increasing importance of clinical nutrition in Turkey and the region, identifying the effects of differences between societies on study results in a clearer way and converting clinical applications into scientific publications as well as forming a bridge between West and East.

The scope of Clinical Science of Nutrition includes original research articles, review articles, case reports, conference reports, and letters to the editor as well as editorials, abstracts from international and national congresses, panel meetings, conferences and symposia. As an online-only publication, in addition to traditional manuscript submissions, Clinical Science of Nutrition is also able to process video, audio and interactive software submissions. Authors, are encouraged to submit their content in the most appropriate medium to best convey their findings to the audience of Clinical Science of Nutrition.

The journal covers all aspects of nutrition and dietetics including prevalence of malnutrition and its effects on clinical results; nutritional support and delivery methods and their advantages and disadvantages; nutritional support products and their side effects; immune system and nutritional support; ERAS protocol and nutritional support; home parenteral and enteral nutrition; nutrition support teams and their necessity, challenges and potential solutions of nutritional support.

EDITORIAL AND PUBLICATION PROCESS

The editorial and publication processes of the journal are shaped in accordance with the guidelines of the International Council of Medical Journal Editors (ICMJE), the World Association of Medical Editors (WAME), the Council of Science Editors (CSE), the Committee on Publication Ethics (COPE), the European Association of Science Editors (EASE), and National Information Standards Organization (NISO). The journal conforms to the Principles of Transparency and Best Practice in Scholarly Publishing (doaj.org/bestpractice).

Originality, high scientific quality, and citation potential are the most important criteria for a manuscript to be accepted for publication. Manuscripts sub-

mitted for evaluation should not have been previously presented or already published in an electronic or printed medium. The journal should be informed of manuscripts that have been submitted to another journal for evaluation and rejected for publication. The submission of previous reviewer reports will expedite the evaluation process. Manuscripts that have been presented in a meeting should be submitted with detailed information on the organization, including the name, date, and location of the organization.

PEER REVIEW PROCESS

Manuscripts submitted to Clinical Science of Nutrition will go through a double-blind peer-review process. Each submission will be reviewed by at least two external, independent peer reviewers who are experts in their fields in order to ensure an unbiased evaluation process. The editorial board will invite an external and independent editor to manage the evaluation processes of manuscripts submitted by editors or by the editorial board members of the journal. The Editor in Chief is the final authority in the decision-making process for all submissions.

ETHICAL PROCEDURES

An approval of research protocols by the Ethics Committee in accordance with international agreements (World Medical Association Declaration of Helsinki “Ethical Principles for Medical Research Involving Human Subjects,” amended in October 2013, www.wma.net) is required for experimental, clinical, and drug studies and for some case reports. If required, ethics committee reports or an equivalent official document will be requested from the authors. For manuscripts concerning experimental research on humans, a statement should be included that shows that written informed consent of patients and volunteers was obtained following a detailed explanation of the procedures that they may undergo. For studies carried out on animals, the measures taken to prevent pain and suffering of the animals should be stated clearly. Information on patient consent, the name of the ethics committee, and the ethics committee approval number should also be stated in the Materials and Methods section of the manuscript. It is the authors’ responsibility to carefully protect the patients’ anonymity.

For photographs that may reveal the identity of the patients, signed releases of the patient or of their legal representative should be enclosed.

PLAGIARISM

Clinical Science of Nutrition is extremely sensitive about plagiarism. All submissions are screened by a similarity detection software (iThenticate by Cross-Check) at any point during the peer-review and/or production process. Even if you are the author of the phrases or sentences, the text should not have unacceptable similarity with the previously published data.

When you are discussing others' (or your own) previous work, please make sure that you cite the material correctly in every instance.

In the event of alleged or suspected research misconduct, e.g., plagiarism, citation manipulation, and data falsification/fabrication, the Editorial Board will follow and act following COPE guidelines.

In the event of alleged or suspected research misconduct, e.g., plagiarism, citation manipulation, and data falsification/fabrication, the Editorial Board will follow and act in accordance with COPE guidelines.

AUTHORSHIP

Each individual listed as an author should fulfil the authorship criteria recommended by the International Committee of Medical Journal Editors

(ICMJE - www.icmje.org). The ICMJE recommends that authorship be based on the following 4 criteria:

1. Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; AND
2. Drafting the work or revising it critically for important intellectual content; AND
3. Final approval of the version to be published; AND
4. Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

In addition to being accountable for the parts of the work he/she has done, an author should be able to identify which co-authors are responsible for specific other parts of the work. In addition, authors should have confidence in the integrity of the contributions of their co-authors.

All those designated as authors should meet all four criteria for authorship, and all who meet the four criteria should be identified as authors. Those who do not meet all four criteria should be acknowledged in the title page of the manuscript.

Clinical Science of Nutrition requires corresponding authors to submit a signed and scanned version of the Copyright Agreement and Acknowledgement of Authorship form (available for download through clinscinutr.org) during the initial submission process in order to act appropriately on authorship rights and to prevent ghost or honorary authorship. If the editorial board suspects a case of "gift authorship," the submission will be rejected without further review. As part of the submission of the manuscript, the corresponding author should also send a short statement declaring that he/she accepts to undertake all the responsibility for authorship during the submission and review stages of the manuscript.

DECLARATION OF INTEREST

Clinical Science of Nutrition requires and encourages the authors and the individuals involved in the evaluation process of submitted manuscripts to disclose any existing or potential conflicts of interests, including financial, consultant, and institutional, that might lead to potential bias or a conflict of interest. Any financial grants or other support received for a submitted study from individuals or institutions should be disclosed to the Editorial Board. To disclose a potential conflict of interest, the ICMJE Potential Conflict of Interest Disclosure Form should be filled in and submitted by all contributing authors. Cases of a potential conflict of interest of the editors, authors, or reviewers are resolved by the journal's Editorial Board within the scope of COPE and ICMJE guidelines.

The Editorial Board of the journal handles all appeal and complaint cases within the scope of COPE guidelines. In such cases, authors should get in direct contact with the editorial office regarding their appeals and complaints. When needed, an ombudsperson may be assigned to resolve cases that cannot be resolved internally. The Editor in Chief is the final authority in the decision-making process for all appeals and complaints.

COPYRIGHT AND LICENSE

Clinical Science of Nutrition requires each submission to be accompanied by a Copyright Agreement and Acknowledgement of Authorship Form (available for download clinscinutr.org). By signing this form, authors retain the copyright of their work and agree that the article, if accepted for publication by the Clinical Science of Nutrition, will be licensed under a Creative Commons Attribution-Non Commercial 4.0 International License (CC-BY-NC).

When using previously published content, including figures, tables, or any other material in both print and electronic formats, authors must obtain permission from the copyright holder. Legal, financial and criminal liabilities in this regard belong to the author(s).

DISCLAIMER

Statements or opinions expressed in the manuscripts published in Clinical Science of Nutrition reflect the views of the author(s) and not the opinions of the editors, the editorial board, or the publisher; the editors, the editorial board, and the publisher disclaim any responsibility or liability for such materials. The final responsibility in regard to the published content rests with the authors.

MANUSCRIPT PREPARATION

The manuscripts should be prepared in accordance with ICMJE-Recommendations for the Conduct, Reporting, Editing, and Publication of Scholarly Work in Medical Journals (updated in December 2018 - <http://www.icmje.org/conflicts-of-interest/>). Authors are required to prepare manuscripts in accordance

with the CONSORT guidelines for randomized research studies, STROBE guidelines for observational original research studies, STARD guidelines for studies on diagnostic accuracy, PRISMA guidelines for systematic reviews and meta-analysis, ARRIVE guidelines for experimental animal studies, and TREND guidelines for non-randomized public behaviour.

Manuscripts can only be submitted through the journal's online manuscript submission and evaluation system, available at clinscinutr.org. Manuscripts submitted via any other medium will not be evaluated.

Manuscripts submitted to the journal will first go through a technical evaluation process where the editorial office staff will ensure that the manuscript has been prepared and submitted in accordance with the journal's guidelines. Submissions that do not conform to the journal's guidelines will be returned to the submitting author with technical correction requests.

Authors are required to submit the following:

- Copyright Agreement and Acknowledgement of Authorship Form and
- ICMJE Potential Conflict of Interest Disclosure Form (should be filled in by all contributing authors)

during the initial submission. These forms are available for download at clinscinutr.org.

Preparation of the Manuscript

Title page: A separate title page should be submitted with all submissions and this page should include:

- The full title of the manuscript as well as a short title (running head) of no more than 50 characters,
- Name(s), affiliations, and highest academic degree(s) of the author(s),
- Grant information and detailed information on the other sources of support,
- Name, address, telephone (including the mobile phone number) and fax numbers, and email address of the corresponding author,
- Acknowledgment of the individuals who contributed to the preparation of the manuscript but who do not fulfil the authorship criteria.

Abstract: An abstract should be submitted with all submissions except for Letters to the Editor. The abstract of Original Articles should be structured with subheadings (Objective, Methods, Results, and Conclusion). Please check Table 1 for word count specifications.

Keywords: Each submission must be accompanied by a minimum of three to a maximum of six keywords for subject indexing at the end of the abstract. The keywords should be listed in full without abbreviations. The keywords should be selected from the National Library of Medicine, Medical Subject Headings database (<https://www.nlm.nih.gov/mesh/MBrowser.html>).

Main Points: All submissions except letters to the editor should be accompanied by 3 to 5 "main points" which should emphasize the most noteworthy results of the study and underline the principle message that is addressed to the reader. This section should be structured as itemized to give a general overview of the article. Since "Main Points" targeting the experts and specialists of the field, each item should be written as plain and straightforward as possible.

Manuscript Types

Original Articles: This is the most important type of article since it provides new information based on original research. The main text of original articles should be structured with Introduction, Methods, Results, and Discussion subheadings. Please check Table 1 for the limitations for Original Articles.

Statistical analysis to support conclusions is usually necessary. Statistical analyses must be conducted in accordance with international statistical reporting standards (Altman DG, Gore SM, Gardner MJ, Pocock SJ. Statistical guidelines for contributors to medical journals. *Br Med J* 1983; 7: 1489-93). Information on statistical analyses should be provided with a separate subheading under the Materials and Methods section and the statistical software that was used during the process must be specified.

Units should be prepared in accordance with the International System of Units (SI).

Editorial Comments: Editorial comments aim to provide a brief critical commentary by reviewers with expertise or with high reputation in the topic of the research article published in the journal. Authors are selected and invited by the journal to provide such comments. Abstract, Keywords, and Tables, Figures, Images, and other media are not included.

Review Articles: Reviews prepared by authors who have extensive knowledge on a particular field and whose scientific background has been translated into a high volume of publications with a high citation potential are welcomed. These authors may even be invited by the journal. Reviews should describe, discuss, and evaluate the current level of knowledge of a topic in clinical practice and should guide future studies. The main text should contain Introduction, Clinical and Research Consequences, and Conclusion sections. Please check Table 1 for the limitations for Review Articles.

Case Reports: There is limited space for case reports in the journal and reports on rare cases or conditions that constitute challenges in diagnosis and treatment, those offering new therapies or revealing knowledge not included in the literature, and interesting and educative case reports are accepted for publication. The text should include Introduction, Case Presentation, and Discussion subheadings. Please check Table 1 for the limitations for Case Reports.

Letters to the Editor: This type of manuscript discusses important parts, overlooked aspects, or lacking parts of a previously published article. Articles on subjects within the scope of the journal that might attract the readers' attention, particularly educative cases, may also be submitted in the form of a "Letter to the Editor." Readers can also present their comments on the published manuscripts in the form of a "Letter to the Editor." Abstract, Keywords, and Tables, Figures, Images, and other media should not be included. The text should be unstructured. The manuscript that is being commented on must be properly cited within this manuscript.

Tables

Tables should be included in the main document, presented after the reference list, and they should be numbered consecutively in the order they are referred to within the main text. A descriptive title must be placed above the tables. Abbreviations used in the tables should be defined below the tables by footnotes (even if they are defined within the main text). Tables should be created using the "insert table" command of the word processing software and they should be arranged clearly to provide easy reading. Data presented in the tables should not be a repetition of the data presented within the main text but should be supporting the main text.

Figures and Figure Legends

Figures, graphics, and photographs should be submitted as separate files (in TIFF or JPEG format) through the submission system. The files should not be embedded in a Word document or the main document. When there are figure subunits, the subunits should not be merged to form a single image. Each subunit should be submitted separately through the submission system. Images should not be

labeled (a, b, c, etc.) to indicate figure subunits. Thick and thin arrows, arrowheads, stars, asterisks, and similar marks can be used on the images to support figure legends. Like the rest of the submission, the figures too should be blind. Any information within the images that may indicate an individual or institution should be blinded. The minimum resolution of each submitted figure should be 300 DPI. To prevent delays in the evaluation process, all submitted figures should be clear in resolution and large in size (minimum dimensions: 100 × 100 mm). Figure legends should be listed at the end of the main document.

All acronyms and abbreviations used in the manuscript should be defined at first use, both in the abstract and in the main text. The abbreviation should be provided in parentheses following the definition.

When a drug, product, hardware, or software program is mentioned within the main text, product information, including the name of the product, the producer of the product, and city and the country of the company (including the state if in USA), should be provided in parentheses in the following format: "Discovery St PET/CT scanner (General Electric, Milwaukee, WI, USA)"

All references, tables, and figures should be referred to within the main text, and they should be numbered consecutively in the order they are referred to within the main text.

Limitations, drawbacks, and the shortcomings of original articles should be mentioned in the Discussion section before the conclusion paragraph.

References

Both in-text citations and the references must be prepared according to the Vancouver style.

While citing publications, preference should be given to the latest, most up-to-date publications. Authors should avoid using references that are older than ten years. The limit for the old reference usage is 15% in the journal. If an ahead-

Type of manuscript	Word limit	Abstract word limit	Reference limit	Table limit	Figure limit
Original Article	5000	300 (Structured)	50	6	7 or total of 15 images
Review Article	6000	300	60	6	10 or total of 20 images
Case Report	2500	250	20	No tables	10 or total of 20 images
Letter to the Editor	1000	No abstract	5	No tables	No media
Editorial	1000	No abstract	5	No tables	No media

of-print publication is cited, the DOI number should be provided. Authors are responsible for the accuracy of references. Journal titles should be abbreviated in accordance with ISO 4 standards. When there are six or fewer authors, all authors should be listed. If there are seven or more authors, the first six authors should be listed followed by "et al." In the main text of the manuscript, references should be cited using Arabic numbers in parentheses. The reference styles for different types of publications are presented in the following examples.

Journal Article: Rankovic A, Rancic N, Jovanovic M, Ivanović M, Gajović O, Lazić Z, et al. Impact of imaging diagnostics on the budget – Are we spending too much? *Vojnosanit Pregl* 2013; 70: 709-11.

Book Section: Suh KN, Keystone JS. Malaria and babesiosis. Gorbach SL, Barlett JG, Blacklow NR, editors. *Infectious Diseases*. Philadelphia: Lippincott Williams; 2004.p.2290-308.

Books with a Single Author: Sweetman SC. *Martindale the Complete Drug Reference*. 34th ed. London: Pharmaceutical Press; 2005.

Editor(s) as Author: Huizing EH, de Groot JAM, editors. *Functional reconstructive nasal surgery*. Stuttgart-New York: Thieme; 2003.

Conference Proceedings: Bengjsson S. Sothemin BG. Enforcement of data protection, privacy and security in medical informatics. In: Lun KC, Degoulet P, Piemme TE, Rienhoff O, editors. *MEDINFO 92. Proceedings of the 7th World Congress on Medical Informatics*; 1992 Sept 6-10; Geneva, Switzerland. Amsterdam: North-Holland; 1992. pp.1561-5.

Scientific or Technical Report: Cusick M, Chew EY, Hoogwerf B, Agrón E, Wu L, Lindley A, et al. Early Treatment Diabetic Retinopathy Study Research Group. Risk factors for renal replacement therapy in the Early Treatment Diabetic Retinopathy Study (ETDRS), Early Treatment Diabetic Retinopathy Study Kidney Int: 2004. Report No: 26.

Thesis: Yılmaz B. Ankara Üniversitesindeki Öğrencilerin Beslenme Durumları, Fiziksel Aktiviteleri ve Beden Kitle İndeksleri Kan Lipidleri Arasındaki İlişkiler. H.Ü. Sağlık Bilimleri Enstitüsü, Doktora Tezi. 2007.

Manuscripts Accepted for Publication, Not Published Yet: Slots J. The microflora of black stain on human primary teeth. *Scand J Dent Res*. 1974.

Epub Ahead of Print Articles: Cai L, Yeh BM, Westphalen AC, Roberts JP, Wang ZJ. Adult living donor liver imaging. *Diagn Interv Radiol*. 2016 Feb 24. doi: 10.5152/dir.2016.15323. [Epub ahead of print].

Manuscripts Published in Electronic Format: Morse SS. Factors in the emergence of infectious diseases. *Emerg Infect Dis* (serial online) 1995 Jan-Mar (cited 1996 June 5): 1(1): (24 screens). Available from: URL: [http:// www.cdc.gov/ncidod/EID/cid.htm](http://www.cdc.gov/ncidod/EID/cid.htm).

REVISIONS

When submitting a revised version of a paper, the author must submit a detailed "Response to the reviewers" that states point by point how each issue raised by the reviewers has been covered and where it can be found (each reviewer's comment, followed by the author's reply and line numbers where the changes have been made) as well as an annotated copy of the main document. Revised manuscripts must be submitted within 30 days from the date of the decision letter. If the revised version of the manuscript is not submitted within the allocated time, the revision option may be canceled. If the submitting author(s) believe that additional time is required, they should request this extension before the initial 30-day period is over.

Accepted manuscripts are copy-edited for grammar, punctuation, and format. Once the publication process of a manuscript is completed, it is published online on the journal's webpage as an ahead-of-print publication before it is included in its scheduled issue. A PDF proof of the accepted manuscript is sent to the corresponding author and their publication approval is requested within 2 days of their receipt of the proof.

Editor in Chief: Sadık Kılıçturgay

Address: Department of General Surgery, Uludağ University School of Medicine, Bursa, Turkey

E-mail: skturgay@gmail.com

Publisher: AVES

Address: Büyükdere Cad. 105/9 34394 Mecidiyeköy, Şişli, İstanbul, Turkey

Phone: +90 212 217 17 00

Fax: +90 212 217 22 92

E-mail: info@avesyayincilik.com

Web page: avesyayincilik.com

CONTENTS

REVIEWS

- 1 **Nursing Practices in Enteral Nutrition**
Kezban Akçay, Derya Suluhan, Gülşah Kesik, Kadriye Uzunoğlu, Esen Kartal, Arzu Şahna
- 15 **Parenteral nutrition and nursing care: A review of the recent literature**
Kezban Akçay, Gülşah Kesik, Derya Suluhan, Derya Soysal, Hacer Nil Demir, Hasibe Kazancı

ORIGINAL ARTICLES

- 21 **The prevalence of obesity in adolescent age group and its relationship with dietary habits, family history, anthropometric measurements and muscle analysis**
Ahmet Bahar, Bahar Özmen Şentürk, Serpil Büyükdemir, Gizem Kutlutürk, Zeynep Nihan Mutlu, Şeyma Baki, Burcu Kulaksız, Mustafa Altinkaynak, Bülent Saka
- 30 **The effect of perioperative nutritional support on readmission ratios in elective major abdominal surgery**
R. Haldun Gündoğdu, Ömer Yazıcıoğlu, Bahadır Osman Bozkırlı, Soner Akbaba
- 35 **Assessment of the STRONGkids screening tool: A cross-sectional study in Turkish children**
Betül Oruçoğlu, Neriman İnanç
- 43 **The trending topic of specialty in Turkey: Clinical nutrition**
Kutay Demirkan, Haldun Gündoğdu, Mehmet Uyar, Sadık Kılıçturgay, Kubilay Demirağ, Osman Abbasoğlu

CASE REPORT

- 49 **Are enteral nutrition solutions effective in patients with diabetic foot wounds?**
Sami Açar

Nursing Practices in Enteral Nutrition

Kezban Akçay¹ , Derya Suluhan² , Gülşah Kesik³ , Kadriye Uzunoğlu⁴ , Esen Kartal⁵ , Arzu Şahna⁶ 

¹Nutrition Support Unit, Hacettepe Adult Hospital, Ankara, Turkey

²Department of Child Health and Diseases Nursing, Health Sciences University Gülhane Faculty of Nursing, Ankara, Turkey

³Department of Internal Diseases Nursing, Hacettepe University Faculty of Nursing, Ankara, Turkey

⁴Nutrition Support Unit, Dr. Abdurrahman Yurtaslan Ankara Oncology Training and Research Hospital, Ankara, Turkey

⁵MH- 1 General Hospital Education Nursing Unit, Ankara State Hospital, Ankara, Turkey

⁶Nutrition Support Unit, Ankara State Hospital, Ankara, Turkey

ORCID iDs of the authors: K.A. 0000-0003-2848-3449; D.S. 0000-0002-7358-7266; G.K. 0000-0002-5748-8225; K.U. 0000-0002-2287-6818; E.K. 0000-0002-4874-184X; A.Ş. 0000-0003-3254-9578.

Cite this article as: Akçay K, Suluhan D, Kesik G, Uzunoğlu K, Kartal E, Şahna A. Nursing Practices in Enteral Nutrition. Clin Sci Nutr 2020; 2(1): 1-14.

ABSTRACT

Enteral nutrition is the preferred route of nutrition when patients have a functioning gastrointestinal tract but are unable to meet their requirements orally. In terms of patient safety and quality of life, it is important that enteral nutrition must be administered by the nutrition team consisting of experienced interdisciplinary health professionals. Members of the nutrition team should have up-to-date knowledge and skills to monitor and evaluate the patients during the enteral feeding process and prevent the development of complications. The nurse, as a member of the team, has important responsibilities, such as correct and safe enteral nutrition practices, care of the enteral nutrition route, and follow-up of patients in accordance with the guidelines.

Keywords: Care, enteral nutrition, nursing

Introduction

Nutrition is an action that needs to be done consciously to get the nutrients needed by the body in sufficient quantity and at the appropriate time in order to maintain the body functions, protect and improve health, and improve the quality of life (1). Malnutrition develops in cases where nutritional therapy cannot be performed or is not sufficiently carried out. Nutritional therapy is needed for the prevention or treatment of malnutrition. Enteral nutrition (EN) is often preferred as the closest method to oral nutrition. EN has many advantages: it is cheap and safe and maintains the structure and functions of the intestine (2).

EN treatment is preferred in patients whose gastrointestinal system (GIS) functions are normal or close to normal and who cannot meet some or all of their daily nutritional requirements (1). EN is used in cases of impaired oral intake for longer than 5–7 days or the prediction of its deterioration, development of malnutrition, unconsciousness, difficulty swallowing, partial intestinal failure, pancreatitis, cancer, mental disorders, cachexia accompanying coma and chronic diseases, metabolic diseases, congenital

heart diseases, digestive system anomalies, when necessary after all types of surgery, and eating disorders (3, 4).

Access routes for enteral nutrition therapy

Nurses must have up-to-date knowledge and skills about EN access routes and nutrition methods for proper and safe administration of EN. The first method to be selected in the patient's diet is oral way. However, in cases where oral feeding is insufficient due to difficulties in swallowing and vomiting, or in cases such as coma where feeding is not possible, EN treatment with nasal/oral tube or gastrostomy tube (GT) is preferred while GIS function is normal (1).

In the selection of the access route for EN treatment, the characteristics of the disease, gastric and intestinal motility and functional status, GI changes due to previous surgery, and the estimated duration of nutritional therapy to be applied are taken into consideration. The most important of these factors is the duration of EN treatment. If nutritional therapy will take less than 4–6 weeks, EN therapy is recommended through nasogastric (NGT)/nasoenteric tube (NET). If the duration is longer than 4–6 weeks, EN

therapy is recommended through enterostomy (gastrostomy or jejunostomy) (2, 4, 5).

Tube feeding is specifically EN provided into the gastrointestinal tract through a tube, catheter, or stoma that delivers nutrients distal to the oral cavity. The tube is inserted through the nose (nasogastric or nasojejunal) or from a stoma (gastrostomy, percutaneous endoscopic gastrostomy with jejunal extension [PEG-J], or percutaneous endoscopic jejunostomy [PEJ]) that opens to the stomach or jejunum endoscopically, surgically or radiologically (1, 4).

Enteral nutrition therapy with nasal tube: NGT and NET are the most common access routes used in the treatment of EN. The most important advantage is that it can be placed easily at the bedside. The direct delivery of foods to the stomach makes it possible to use hypertonic foods and to deliver them at higher speeds and in bolus style (4, 6, 7). NGTs are used in adult patients. NGTs used for gastric access are preferred for less than 4–6 weeks (Table 1). The lengths of the tubes range from 90 to 120 cm; their diameter varies between 6 and 22 French; and they have one/three lumen tube types made of more rigid polyvinyl, more flexible silicone, or less traumatic polyurethane material (7, 8).

Evaluation of whether NGT or NET is in place before every use is an important nursing intervention in terms of safe application and patient safety. Radiological evaluation (x-ray), the measurement of the distance between the proximal end of the tube and the nose, and pH control are the methods used to evaluate the placement of the tube. Radiological evaluation is recommended as the gold standard in the assessment of tube placement in the guidelines of the American Society for Parenteral and Enteral Nutrition (ASPEN) (2009) and the Dietitians Association of Australia (2018) (7, 9). The American Association of Inten-

sive Care Nurses suggests that the tube location should be marked with the indelible pen at the edge of the nose, and this mark should be checked every four hours and before each feeding/drug application (10). If the outer length of the tube is stable and the pH of gastric aspirate is less than 5, NGT is placed in the stomach (8). In the treatment of nutrition with the nasojejunal tube, double lumen gastrojejunal tubes may be preferred in patients who require gastric decompression with small bowel nutrition. These tubes are placed by experienced physicians under the guidance of endoscopy or fluoroscopy (6, 7).

Nutrition therapy with gastrostomy tube: Nutrition with the GT is a fairly common and safe procedure. For long-term EN treatment (more than 4–6 weeks), if delayed gastric emptying and obstruction are not present, nutritional therapy with GT is preferred (2, 4). EN treatment with GT is considered superior to other methods for the following reasons: gastric acid having a bactericidal effect, normal functions of pancreatic enzymes, stomach being more appropriate for bolus nutrition, better tolerance of hyperosmolar nutrients, reduced frequency of lung infections, and its ease of application and comfort level (4). In cases where gastrostomy is contraindicated, jejunostomy is the preferred access route for EN therapy in patients with a risk of aspiration pneumonia (8).

Oral nutritional supplements

Oral nutritional therapy is essentially the delivery of oral nutritional supplements and defined as “foods with special medical purpose” (1). Oral nutritional supplements are the solutions to meet macro- and micronutrient needs. They are ready-made liquids, creams, or powdered supplements that can be added to drinks or food. Liquid oral nutritional supplements that are ready for drinking or made from powders are sometimes referred to as sip feeds (6).

Table 1. Features of tubes according to their types (9)

Features	Polyvinyl chloride (PVC) tubes	Polyurethane (PE) and silicone tubes
Measurement (French)	6–12 Fr	14–22 Fr
Purpose of usage	Gastric drainage, gastric aspiration, and gastric decompression	Enteral nutrition
Structure	Hard Soluble in stomach contents	Soft
Patient comfort	Comfortless	Comfortable
Approximate time of use	10 days	4–6 weeks
Wire guide	None	Yes/none
Type	Levin tube Ryles tube Salem tube	Many types

Enteral Nutrition Application Methods

EN products are given to the digestive system that becomes functional by using one of the continuous or intermittent nutrition methods.

Continuous nutrition: The nutritional solution is given to the patient continuously by using a pump without interruption (16–24 hours). With this method, the EN product is given at a constant speed with the help of a feeding pump throughout the day or at a desired time continuously (4).

It has advantages such as reducing the risk of aspiration resulting from reflux or high gastric residual volume (GRV), providing controlled delivery into the intestine, and help in controlling blood glucose level (4, 8, 11, 12).

Intermittent nutrition: When nutrition is carried out at the decided time in intermittent or cyclic nutrition method (one hour or more), the nutrition is stopped for four to eight hours during the rest of the day, and the gastrointestinal system is rested. In intermittent feeding, EN treatment is carried out using the hanging feeding bag or a feeding injector with the effect of gravity. Intermittent feeding can be applied three to five times a day by bolus or cyclically. While bolus nutrition can be applied with NGT or GT, its application with the jejunal tube is contraindicated.

In bolus feeding method, the plunger of the syringe is removed during EN treatment. The injector body is attached to the distal end of the tube to be used as a funnel through which the enteral product is provided with the effect of gravity. It has many advantages such as providing fast treatment, not requiring an infusion pump, and allowing the stomach to rest between feeding periods (12, 13). Although there is no significant difference between continuous and intermittent nutrition, continuous nutrition is the preferred choice because of the reduction of GI complications and ease of applying EN treatment (8, 14, 15).

Nursing practices in Enteral nutrition

It is important to perform nursing practices in patients receiving EN in line with the recommendations of the guidelines that provide up-to-date and high-level information in order to ensure patient safety and increase the quality of life. The nurses' responsibilities in the safe delivery of EN therapy include effective interdisciplinary communication, EN implementation, maintenance of delivery routes, and patient monitoring and recording (16-21).

Application of enteral nutrition therapy: Enteral products are given to the patient with NGT in short-term nu-

tritional support and via enterostomy (gastrostomy, jejunostomy) in long-term nutritional therapy (2, 4, 5). In our country, on 19 April 2011, the Ministry of Health published the "Regulation on the Amendment of the Nursing Regulation." According to this regulation, it is the responsibility of the nurse to place and remove the NGT with the decision of the physician (22). Nursing practices in EN are included in Tables 2, 3, and 4 within the scope of this responsibility.

One of the issues to be considered in the implementation of EN therapy is the correct understanding of the preapplication treatment protocol. In the ASPEN (2017) guideline, it is recommended to check the patient's information, the enteral feeding route (NET, NGT, GT, PEG-J), the EN method (intermittent/continuous/bolus feeding), and the feeding speed before giving EN to the patient (8).

Maintenance of access routes: Maintenance of access routes includes dressing of the enteral tube and maintaining the patency of the path. The dressing of the enteral tube varies according to the type of tube the patient has. In NGT maintenance, the tapes used for tube fixing should be changed daily. Before replacing the fixed tape with a new one, the distal length of the tube should be checked to determine if the tube is in place. To prevent trauma that may occur in the nasal mucosa, the tape should be fixed to another side of the nostril at each time (18). The nasal mucosa should be observed daily for any irritation (18, 19). In case of irritation in the nasal mucosa, the tube should be removed and inserted through the other nostril and hypoallergic and disposable products that ensure secure fixation should be preferred (18). The maintenance of the peristomal area (GT, PEG-J) is performed using the aseptic technique to reduce the risk of local infection, according to local wound management policy. Although the use of dressings is not recommended, the European Society of Clinical Nutrition and Metabolism (ESPEN) recommends a sterile GT dressing the next morning and the first 7 days following the placement of the PEG tube (10). In the first 48 hours after the insertion of the GT, the peristomal area should be evaluated for bleeding (a small amount of bleeding may occur) and hematoma.

The peristomal area should be evaluated at 4-hour intervals in the first 24 hours following the insertion of gastrostomy with the monitoring of vital signs and then should be followed the clinical procedure. In the evaluation of peristomal area, symptoms such as redness, increased temperature, increased pain, excessive discharge, or odor may show local infection (peristomal infection). The presence of leakage (contents of the stomach) from the edge of the tube may indicate that the location of the tube has changed.

Table 2. Nursing interventions and reasons in the application of enteral nutrition therapy with nasogastric tube (8, 17-22)	
Interventions	Reasons
The individual's condition and medical records are evaluated: Nasal patency, swallowing reflex, and bowel movements	If bowel sounds are not heard, peristaltic movements may not have returned. The risk of aspiration and abdominal distention increases.
Order is checked.	The correct application is realized.
Materials to be used are prepared.	<ul style="list-style-type: none"> • Enteral formula (at room temperature) • Emesis basin • Towel or paper towel • Container for dirty objects • Disposable gloves • 50-mm feeding injector/feeding bag • Water at room temperature
All materials are brought to the individual, in an area that you can easily reach.	The process is made easier and saves time.
The process and its purpose are explained to the individual in detail and his/her approval is obtained.	This application relieves the anxiety of the individual. It ensures compliance and participation.
Hands are washed and gloves are worn.	You prevent the spread of microorganisms in your hands.
Unless there is a medical indication, the patient's head is raised at least 30–45 degrees in lying position.	It decreases the risk of aspiration.
Towels or paper towels are laid on the chest of the individual.	The contamination of the individual's clothes is prevented.
The tip of the nasogastric tube is clamped, and the connections of the tube are removed.	Leakage of stomach contents or entry of air into the stomach is prevented.
After placing the injector at the end of the tube, the clamp on the tube is opened and, if necessary, the contents of the stomach are drawn into the injector by withdrawing the plunger of the injector.	It gives information about whether previously given enteral formula is tolerated by the digestive system.
<p>If the enteral formula is to be given with a feeding bag: The enteral formula in the box/bottle at room temperature is shaken and filled into the feeding bag, and the bag is hung on the serum rack. The formula poured into the bag is passed through the set, and the set air is removed. Before enteral nutrition therapy, the recommended amount of water at room temperature is given to the patient.</p> <p>If the enteral formula is to be given with a feeding injector: The recommended amount of water at room temperature before feeding is given to the patient through the tube with an injector. The plunger of the feeding injector is removed, and the injector body is placed at the distal end of the probe to be used as a funnel. The tube is clamped with the help of hand during the addition of the formula to the injector. The enteral formula in the box/bottle at room temperature is filled into the injector by shaking. The nutritional injector is kept 30–35 cm above the level of the stomach so that it can reach there with the effect of gravity. The flow rate is adjusted by the height of the injector.</p>	The entry of air into the stomach is prevented.

Table 2. Nursing interventions and reasons in the application of enteral nutrition therapy with nasogastric tube (8, 17-22) (Continued)

Interventions	Reasons
After the enteral nutrition treatment is finished, the recommended amount of water at room temperature is given through the tube.	The formula left in the tube is allowed to go from the tube to the stomach, and the tube is prevented from clogging.
When enteral nutrition therapy is finished, the cap at the distal end of the tube is closed.	The formula may leak out.
Oral care is given if necessary.	The individual is provided with care to feel comfortable.
Tools and equipment are properly removed from the environment.	The spread of microorganisms will be prevented.
If contraindicated, for 30–60 minutes after enteral nutrition treatment, the head of the individual is left at 30–45 degrees.	It is necessary for the prevention of aspiration.
Gloves are removed and hands are washed.	The spread of microorganisms will be prevented.
The application, the individual's response, and participation are recorded.	It provides data for the individual's subsequent procedures and provides legal basis for the nurse.

Maintaining the opening of the access route and tube maintenance: Despite the differences in the amount of water used and the time of delivery of water in maintaining the opening of the access route in clinical practice, in the ASPEN guide (2017), it is recommended that feeding tube should be washed with minimum 30-mL water every four hours during continuous nutrition and before and after feeding in intermittent nutrition (8).

The tube should be regularly washed with water before and after drug administration during intermittent or continuous feeding in order to prevent the complication of tube obstruction (23). In cases where EN treatment with GT is not allowed, the dysfunctional GT should be changed in line with the institutional policies and manufacturer's recommendations. In the ASPEN (2017) guideline, recommendations regarding the replacement of tubes are given below (8):

1. Development of institutional protocols including manufacturer's instructions for changing enterotomy tubes.
 - a. Routine removal and replacement of a well-preserved PEG may not be necessary.
 - b. It is recommended to change it according to the manufacturer's instructions.
2. If any of the following is identified in the patient, the tube replacement may be considered earlier than the time specified in the manufacturer's manual.
 - a. Nonfunctional tube.
 - b. Balloon rupture in tubes, internal fixator of which is balloon.

- c. Presence of peristomal infection that persists despite appropriate antimicrobial therapy (local infection).
 - d. Skin maceration.
 - e. Ulcer formation that does not heal despite good wound care practice.
 - f. Colocutaneous or gastrocolic fistulas.
3. It is recommended that percutaneous tube should be changed after stoma maturation (30–90 days after the initial placement of the tube) or according to institutional protocols.
 4. Routine replacement of the percutaneous tube after stoma maturation (30 days after the initial placement of the tube) or in accordance with institutional protocols may be considered.

Ensuring tube safety: Because bulky dressings between external fixators and the skin can increase tube tension, they can lead to displacement of the GT or injury. It is also necessary to make sure that the external fixator is not too tight or too loose to prevent embedded buffer syndrome and complications such as hypergranulation. In the literature, it is recommended to check the distance between the tube's external fixator and the skin while the patient is in a sitting position, and the ideal distance should be 2–5 mm (10, 13, 20, 21).

Tube rotation: In seamless tubes, unless contraindicated, standard GT (with or without balloons) should be rotated 360 degrees per day. Rotating the tube reduces physical pressure on the gastric mucosa. It is recommended to start rotating the GT 24 hours after the first insertion and to do it daily (10, 13).

Table 3. Nursing interventions and reasons in the application of enteral nutrition with gastrostomy/ jejunostomy tube (8, 17-21, 30)	
Interventions	Reasons
The individual's condition and medical records are evaluated: Ostomy region, gastrostomy/jejunostomy tube and bowel movements.	It gives information about the patient's peristomal area and gastrointestinal system.
Order is checked.	The correct application is deployed.
Materials to be used are prepared.	<ul style="list-style-type: none"> • Connection set (only required for button gastrostomy tube) • Enteral formula • Emesis basin • Towel or paper towel • Container for dirty objects • Disposable gloves • 20–50 mm feeding injector • Water at room temperature
All materials are brought to the individual, in an area that you can easily reach.	The process is made easier and saves time.
The process and its purpose are explained to the individual in detail and his/her approval is obtained.	This application relieves the anxiety of the individual. It ensures compliance and participation.
Hands are washed and gloves are worn.	You prevent the spread of microorganisms with your hands.
Unless contraindicated, the patient's head is raised at least 30–45 degrees in lying position.	It decreases the risk of aspiration.
After placing the injector at the end of the gastrostomy tube, the clamp on the tube is opened, and, if necessary, the contents of the stomach are drawn into the injector by withdrawing the plunger of the injector.	It gives information about the tolerance of the enteral formula previously given by the digestive system.
The recommended amount of water is given from the gastrostomy tube before enteral nutrition therapy.	The opening of the tube is provided.
If the enteral formula is to be given with a feeding bag: After shaking the feeding formula at room temperature, it is poured into the feeding bag, and the bag is hung on the serum rack. Air in the feeding set is discharged with enteral formula.	The nutrition formula at room temperature prevents abdominal distension and cramp. Shaking the formula ensures that the product is poured homogeneously into the bag.
The tip of the feeding set, with the air removed within it, is placed in the PEG or gastrostomy tube. If the patient has a buttoned gastrostomy tube, the feeding set added to the tip of the feeding bag is carefully placed in the buttoned gastrostomy tube.	The feeding set provides a connection between the feeding bag and buttoned gastrostomy tube.
Maximum amount of total product to be given with 6–8 hours should be added to the feeding bag. The speed of the enteral formula, to be sent hourly, is adjusted in the feeding machine. Nutritional product is given to the patient.	Enteral products opened at room temperature should be consumed within a maximum of 6–8 hours. (The maximum time for keeping the formulas at room temperature may differ from each other. Therefore, the instructions for use of the preferred formula should be carefully examined.)
If the enteral formula is to be given with an injector:	
Remove the plunger from the syringe and place the tip of the syringe into tube connector at end of the PEG or gastrostomy tube	

Table 3. Nursing interventions and reasons in the application of enteral nutrition with gastrostomy/ jejunostomy tube (8, 17-21, 30) (Continued)

Interventions	Reasons
If the patient has a button gastrostomy feeding tube (low profile gastrostomy feeding tubes), a water-filled extension set is carefully placed in a buttoned gastrostomy tube. The plunger of the feeding injector is then removed, and the injector body is placed at the distal end of the feeding set to be used as a funnel.	The feeding set acts as a connection between the feeding bag and buttoned gastrostomy.
When enteral nutrition therapy is completed, the recommended amount of water is given through the gastrostomy tube.	Blockage of the feeding probe is prevented.
When the procedure is completed, the cap at the distal end of the probe is closed.	Food can leak out.
If it is not contraindicated, it is ensured that the head of the bed remains elevated at 30–45 degrees at least for 30–60 minutes after enteral nutrition therapy.	The development of aspiration is prevented.
Oral care is given if necessary.	The individual is provided oral care to feel comfortable.
Tools and equipment are properly removed from the environment.	
Gloves are removed and hands are washed.	The spread of microorganisms will be prevented.
Information about application of enteral nutrition, formula type, and amount are recorded.	It provides data for the individual's subsequent procedures and provides legal basis for the nurse.

Tube position: The centimeter mark of the tube at the peristomal skin level (except for low-profile GT) should be recorded and observed before each use (24). A deviation from the first recorded measurement may indicate the movement of the GT. Thus, to prevent any complication, the nurse should inform the physician about the change in the position of the tube before using it.

Access route-related complications and nursing interventions in enteral nutrition

Once EN therapy is initiated, minor and major complications can sometimes occur. Minor complications of EN with GT include pulmonary aspiration, peritonitis, gastrocolic fistula, bleeding, necrotizing fasciitis, and embedded buffer syndrome. Major complications include occlusion of the tube caused by the dysfunctional tube, the fracture or rupture of the tube, the displacement of the tube, dislocation of the tube, and leakage around the tube. In the EN guideline of ASPEN (2017), it is emphasized that nursing care and discharge training are important in preventing minor complications related to GT and enterostomy (8). Nurses should provide training to caregivers or patients about the causes of complications and the prevention methods, to apply appropriate nursing interventions in the treatment of complications, to evaluate and monitor the patient for safe EN treatment. The reasons and solutions for common

complications are given in Table 5. Although there are no research results that provide very strong evidence for the prevention and treatment of some complications, recommendations for some complications in the table are feasible interventions in line with clinical procedures.

Complications related to enteral nutrition and nursing interventions

Diarrhea, nausea or vomiting, regurgitation or aspiration, and refeeding syndrome are common complications of EN therapy. Knowing and preventing the causes and proper treatment of these complications are the responsibilities of the nurses of the nutrition team. Table 6 contains the causes of common EN-related complications and applications for their prevention and treatment.

One of the common complications associated with EN is aspiration (Table 6). Oropharyngeal secretions and/or reflux of stomach contents into the esophagus may be associated with aspiration. Intensive care patients and patients with swallowing disorders may have problems with respiratory tract (8). The American Association of Critical-Care Nurses (AACN) recommends to place the head of the bed at 30–45 degrees or higher for 30–60 minutes during or after the completion of EN, if not contraindicated to reduce the risk of aspiration (25).

Table 4. Nursing interventions and reasons in drug applications from enteral feeding tube (8, 14, 15, 17)	
Interventions	Reasons
Medicine cards written according to the physician's order are checked.	It prevents wrong applications, and it is determined whether the drug order is changed. Thus, errors that may occur are prevented.
Information about the effect and side effect of the drug to be applied is obtained.	This information helps to evaluate the therapeutic effects of drugs.
Materials to be used are prepared.	<ul style="list-style-type: none"> • Drug card • Drug to be used • Drug glass • Drug tray • Injector/scale (liquid drugs) • Drug divider (solid drugs) • Water/fruit juice
Hands are washed.	It prevents the spread of microorganisms.
The name of the drug, its dosage, the form and time of administration, and the patient's name and surname are checked again from the drug card.	It prevents the administration of missing drug, missing dose, or overdose of drug.
When necessary, clinical pharmacist is consulted for the suitability of dosage forms, prevention of interactions, and administration of medication from the feeding tube.	It ensures safe drug administration.
The drug is properly prepared.	
Preparation of liquid dosage forms:	
The preparation is shaken properly before use.	It provides accurate dosing.
Medicines in liquid form are diluted according to the pharmacist's recommendation.	It prevents clogging of the tube.
Medicines in liquid form should not be added directly to the enteral nutrition formula.	It prevents interaction of drug with enteral formula.
Preparation of solid dosage forms:	
The tablets are simply compacted and ground, and the powder obtained is mixed with sterile water.	
If necessary, tablets are crushed to a fine powder in the mortar. 5 mL of water is added, and crushing is continued with a pestle. 5 or 10 mL of water is added again with continuous crushing and mixing, thus creating a good suspension. The suspension is drawn into the syringe of appropriate type and size and applied through the tube.	
In order to avoid any medication in the mortar, 10–20 mL of water is added to the mortar, and it is drawn with the syringe and then applied from the tube. If necessary, this process is repeated until there is no medicine left in the mortar.*	
Effervescent tablets are diluted with 1/2–1/3 glass of water.	
Hard gelatin capsules containing powdered medicines are opened according to the pharmacist's recommendation and diluted with 10–15 mL of water to form a slurry.	
Capsules containing granules should not be crushed after opening.	
In the application of the prolonged-release capsule containing enteric-coated pancreatic enzymes from the feeding tube according to the pharmacist's recommendation, the content is mixed with apple juice.	

Table 4. Nursing interventions and reasons in drug applications from enteral feeding tube (8, 14, 15, 17) (Continued)

Interventions	Reasons
Hard gelatin capsules are opened according to the pharmacist's recommendation, and the powder is mixed with sterile water.	
Soft gelatin capsules are applied to the feeding tube after being pierced with a needle and the liquid contents are drawn with the injector.	
According to the pharmacist's recommendation, the capsule can be put into warm water, and the whole mixture can be applied from the tube.	
Drugs are applied separately.	
Drug should not be added directly to the enteral nutrition formula.	
Before applying the drug, feeding is stopped and, if not contraindicated, the tube is washed with at least 15 mL of sterile water.	Drug interaction is prevented with enteral formula.
Oral injectors with the instruction "only for oral use" are used for drug administration from the feeding tube.	Drug interaction is prevented with enteral formula.
To avoid altering the bioavailability of the drug, enteral nutrition therapy may be delayed for 30 minutes or more, if appropriate.	Drug interaction is prevented with enteral formula.
After applying all medicines, if not contraindicated, the tube is washed once more with 30 mL of water.	Drug interaction is prevented with enteral formula.
Information on drug administration, form, amount of drug, application time, and the person performing the application are recorded.	It provides data for the individual's subsequent procedures and provides legal basis for the nurse.

*This method should be used carefully for patients with fluid restriction.

- Unless contraindicated, the head of the bed is 30–45 degrees.
- Reduce sedative drug use as much as possible.
- Evaluation of the location of the feeding tube at four-hour intervals.
- Observing the external length of the tube.
- Evaluation of GI intolerance (nausea, abdominal distension, and pain) at four-hour intervals.
- Avoid bolus feeding if there is a high aspiration risk.
- Evaluation of swallowing before starting oral feeding for patients extubated after prolonged intubation.
- Keeping the endotracheal tube cuff pressure at an appropriate level.
- Removal of secretions on the cuff before reducing the endotracheal tube cuff pressure.

Current data on gastric residue volume measurement:

One of the factors indicating the nutritional tolerance of the patient receiving EN therapy is GRV. How to measure GRV, when to measure, and what a high GRV actually implies are the major controversies for the clinicians. Many variables, including the type of feeding tube and the patient position, can affect GRV measurements. The control of GRVs in patients with GT in the intensive care setting varies according to local hospital policy. There is no evidence in the literature to suggest the control of GRVs in patients with GT out of the setting of the intensive care. While some studies suggest GRV control (26–28), some do

not recommend it (29). The acceptable range for GRV can vary significantly. The acceptable range for GRV has not been established in studies conducted to date. However, the British Association for Parenteral and Enteral Nutrition (BAPEN) (2003) recommends that the stomach should be aspirated every four hours in patients with suspected gastrointestinal motility, and a nutritional policy should be revised if the gastric aspirates exceed 200 mL (Table 7) (30).

Patient monitoring: Monitoring of the patient with EN by an experienced nutrition support team contributes to the prevention of complications (31). Objectives of patient monitoring are to reach the target body weight and clinical laboratory findings in the normal range and to improve the quality of life (8, 31). ASPEN (2017) and BAPEN (2003) suggest performing the following applications in the follow-up of the patient with EN (8, 30).

- Monitoring and evaluating the patient with EN treatment to determine physical examination findings, laboratory values, anthropometric data, and all changes in parameters (8).
- Evaluation of the nutritional risk during the treatment of the patient with EN (8):
 - a. Determining the frequency of reevaluation considering the clinical course of the patient.
 - b. Establishing protocols related to patient assessment on a daily and/or weekly basis, monitoring nutritional status, and giving more attention to data on reevaluation.

Table 5. Complications related to gastrostomy (8, 10, 13, 19-21, 30-32)	
Complication	Suggestions
Tube-related complication Tube blockage Reason <ul style="list-style-type: none"> • Unproper usage of the tube (use of drugs with high viscosity, small tube lumen, use of gravity method instead of feeding machine for EN) 	<ul style="list-style-type: none"> • Preference of liquid forms of medicines, if possible • Washing the tube before and after each feeding/drug application • Washing the feeding tubes with 30 mL of water every 4 hours during continuous feeding or before and after intermittent feeding in an adult patient (Evidence A) • Washing the feeding tube with 30 mL of water after GRV measurements in an adult patient (Evidence B) • Washing of feeding tubes in newborns and pediatric patients with the minimum volume required to clean the tube (Evidence C) • It is recommended to use sterile water before and after drug administration in adult and newborn/pediatric patients (Evidence C) • Following protocols of proper washing of the tubes before and after drug administration (Evidence B) • Using sterile water for washing of the tubes in patients with weakened immunity or critical illness due to lack of evidence suggesting the use of tap water (Evidence C)
Tube displacement Reason <ul style="list-style-type: none"> • Accidental removal of the tube • In gastrostomies with an internal fixator balloon, the bursting of the balloon or decreasing the amount of balloon fluid used to inflate the balloon • Tube moves toward the intestine 	<ul style="list-style-type: none"> • Fixing the external extension to the skin with tape • Preference of tubes with external fixator • Recording of tube type and external length • Avoiding using tubes that are not intended for long-term enteral feeding, such as urine or gastrointestinal drainage tubes without external fixator • Checking the tube balloon • Avoiding the use of the tube until it is confirmed to be in the right place
Embedded buffer syndrome Reason <ul style="list-style-type: none"> • The tube is not rotated • External fixative is too loose or too tight 	<ul style="list-style-type: none"> • The tube is pushed about 2–3 cm every day and carefully retracted until the inner buffer resistance is felt • Rotation of the tube to 180–360 degrees around its axis (rotation of jejunal or gastrojejunal tubes is not recommended) • The distance between the external fixator and the skin is 2–5 mm
Stoma-related complication Leak around the tube Reason <ul style="list-style-type: none"> • Cases in which the pressure in the abdomen increases (constipation, cough, ventilated patient, vomiting, etc.) • Bursting of balloon in balloon gastrostomy tubes/reduced amount of balloon fluid used to inflate the balloon • The size of the tube (French) not suitable for the stoma entry point • Tube displacement • Nutrition intolerance 	<ul style="list-style-type: none"> • Replacing the tube • Balloon volume control in balloon tubes • Inflation of the balloon • To verify that the tube is not displaced, it is necessary to check the numerical mark on the tube in the stoma outlet area every 4 hours or before using the tube. • Confirmation of tube placement in adults by radiography if the tube is suspected to be displaced • Using barrier products • Dressing and closing with gauze/or wound care materials

Table 5. Complications related to gastrostomy (8, 10, 13, 19-21, 30-32) (Continued)

Complication Tube-related complication	Suggestions
Local irritation Reason <ul style="list-style-type: none"> • The contact of gastric content to the skin • Harsh cleansing of peristomal skin • Use of antibacterial or other topical drugs 	<ul style="list-style-type: none"> • Using barrier products to protect and heal the skin • Use of absorbable dressing material
Local infection Reason <ul style="list-style-type: none"> • Poor hygiene • MRSA 	<ul style="list-style-type: none"> • Cleaning 2–3 times a day with sterile water or physiological serum • Avoid excessive moistening of the peristomal area • Taking culture • Use of ordered antibiotics • Use of silver dressing
Hypergranulation tissue <ul style="list-style-type: none"> • Reason • Excessive mobility of the tube in the stoma • Excessive moisture • Use of tube with inadequate size 	<ul style="list-style-type: none"> • Ensuring and maintaining the tube's stability • Fixing the external extension of the tube to the skin with tape • To prevent excess moisture that may collect in the peristomal region, not covering the stoma with dressing materials (gauze, wound care products), and leaving the peristomal area open. • Using appropriate-sized (French) tube for the stoma • Short-term use of ordered corticosteroid creams (triamcinolone acetonide 5%) (3 times a day, 5–7 days) • Silver nitrate usage: up to 5 days or once a day until hypergranulation tissue shrinks. • Using barrier powder or foam dressing to control moisture

EN: enteral nutrition; GRV: gastric residual volume; MRSA: methicillin-resistant *Staphylococcus aureus*

- Daily monitoring of fluid balance (30).
- Evaluation of body weight or body mass index twice a week or more frequently if there is concern of hydration (30).
- If the patient is receiving EN treatment in the acute unit, daily evaluation of vital signs such as body temperature, pulse, and breathing to detect signs of infection and dehydration (30).
- Daily evaluation of the intestine to determine EN tolerance (30).
- In order to detect hypo-/hyperglycemia, daily evaluation of capillary blood glucose at the beginning until stability is achieved, and once in every four hours daily if the patient has diabetes (30).
- Daily oral care (30).
- Daily evaluation of the physical structure, position, and peristomal area of the tube (30).
- Checking the GT rotation daily, and checking the balloon volume in ballooned GTs weekly (30).
- Evaluation of sodium, urea, and creatinine levels daily until stability is achieved in line with clinical procedures. In case of refeeding syndrome, evaluating them daily, three times a week until stabilized, then as clinically indicated (30).
- Assessment of C-reactive protein twice a week until stability is achieved, assessment of the acute phase response, and interpretation of protein and micronutrient results.
- Weekly evaluation of albumin until stabilized, since low albumin reflects disease rather than protein status (30).
- Assessment of complete blood count twice a week until stabilized to check on infection and anemia (30).
- Evaluation of zinc, copper, selenium, folate, and vitamin B12 if there are clinical indications (30).
- Evaluation of vitamin D in patients with six-month long-term EN or generally in patients receiving EN treatment at home (30).
- Reevaluation of the patient who has received long-term ET nutrition in the hospital at least once a month (8).
- Reassessment of a patient having received ET nutrition at home for a long-term at least once in every three months (8).
- Recording the care and treatment of patient with EN ensures effective communication with the nutrition support team and health care professionals between clinical units or shifts (16).

Table 6. Complications seen during enteral nutrition (13)	
Complication	Prevention/treatment
Diarrhea Reason <ul style="list-style-type: none"> • High infusion rate • Bolus feeding intolerance • Nutrition with high osmolarity • Microbial contamination of enteral product 	<ul style="list-style-type: none"> • Switching to hydrolyzed formula or modular feeding • Decreasing the feeding rate and increasing it as tolerated • Delivering frequent, smaller amounts of food or switching to continuous feeding • Use of sterile, commercially manufactured products if possible • Preparation of nutritional product in clean environment
Nausea or vomiting Reason <ul style="list-style-type: none"> • Drugs (antibiotic, laxative, etc.) • High infusion rate • Delivery of drug during feeding • Psychological factors 	<ul style="list-style-type: none"> • Checking the drugs • Decreasing infusion rate • Giving prokinetics • Gaining regular intestinal habit by using sufficient liquid, fibrous foods, and/or laxatives • Interruption of continuous feeding during drug administration, not performing nutrition and drug administration at the same time • Review of nutritional behavior; considering referral to a psychologist
Regurgitation or aspiration Reason <ul style="list-style-type: none"> • Gastroesophageal reflux • Clogged tube • High infusion rate • Intolerance of bolus feeding 	<ul style="list-style-type: none"> • Correct position, enteral product thickener, drugs, continuous nutrition, jejunal tube, fundoplication • Checking the tube regularly • Decreasing infusion rate • More frequent feeding with less amount or continuous infusion
Refeeding syndrome Reason	<ul style="list-style-type: none"> • Identification of patients at risk of refeeding syndrome before starting EN • Risk factors include: <ul style="list-style-type: none"> • Malnutrition for more than 2 weeks • Poorly controlled diabetes • Before and during cancer treatment • Anorexia nervosa • Short bowel syndrome • Inflammatory bowel disease • Being an elderly patient living alone • Low birth weight and preterm birth • Chronic infections • Monitoring of fluid balance, daily body weight and electrolyte status (potassium, magnesium, and phosphorus), and other metabolic parameters (glucose) depending on the clinical condition of the patient • Evaluation of metabolic and nutritional parameters before starting enteral nutrition therapy • Starting 25% of target requirements on the first day of enteral nutrition therapy • Delivery of complementary thiamine (intravenously or orally) when enteral nutrition is started • When starting enteral nutrition, monitoring parameters such as serum potassium, phosphorus, magnesium, and glucose

EN: enteral nutrition

Conclusion

One of the responsibilities of the nutrition support team is to have the knowledge and skills to monitor and evaluate the patient during the EN treatment and to prevent the development of complications. The nurse,

who is a member of the team, has important responsibilities in safe EN implementation and patient monitoring. Nurses' performance of nutritional practices in accordance with current literature and the contribution of evidence-based research results to practice are important in preventing complications associated with

Table 7. Current recommendations regarding measurement of gastric residual volume (30)**Suggestions for application**

- Evaluate all patients fed with tube in terms of risk of aspiration (Evidence A)*.
- Make sure that the feeding tube is in the correct position before you start feeding (Evidence A)*.
- Always keep the head of the bed 30–45 degrees high during enteral feeding (Evidence A)*.
- For patients with gastric feeding, check GRVs every 4 hours for the first 48 hours. After the EN target rate is reached, patients who are not in the intensive care can be monitored for GRVs in every 6–8 hours (Evidence C)*.
- In intensive care patients, measurement can be done every 4 hours (Evidence B)*.
- In adult patients, a promotility agent should be considered if GRV is 250 mL or more after the second measurement (Evidence A)*.
- If GRV is more than 500 mL, EN treatment should be stopped using an algorithm for physical evaluation, gastrointestinal evaluation, evaluation of glycemic control, minimizing sedation, and patient's nutritional tolerance (Evidence B)*.
- In sick children who constantly receive EN, GRVs can be checked every 4 hours, and EN is stopped if GRV is higher than or equal to the hourly rate. If the child receives EN by bolus method, GRV can be checked before the next feeding, and EN is stopped if GRV is more than half of the previous feeding volume (Evidence C)*.

EN: enteral nutrition; GRV: gastric residual volume. *Evidence definitions in ASPEN 2017 guideline (8): Evidence A = There is good research-based evidence to support the guideline (prospective, randomized studies). Evidence B = There is appropriate research-based evidence to support the guideline (well-designed studies without randomization). Evidence C = Based on guideline, expert opinion, and editorial consensus

EN, reducing the length of hospital stay, and increasing the quality of life.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept – D.S., K.A.; Design – D.S., K.A.; Supervision – A.Ş., E.K., K.U.; Resources – D.S., K.A., E.K.; Analiz ve/veya Yorum / Analysis and/or Interpretation – D.S., K.A., K.U., G.K., A.Ş., E.K.; Literature Search – D.S., G.K.; Writing Manuscript – D.S., K.A.; Critical Review – D.S., K.A., K.U., G.K., A.Ş., E.K.

Conflict of Interest: The authors have no conflicts of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

References

1. Cederholm T, Barazzoni R, Austin P, Ballmer P, Biolo G, Bischoff SC, et al. ESPEN guidelines on definitions and terminology of clinical nutrition. *Clin Nutr* 2017; 36: 49-64. [\[Crossref\]](#)
2. ASPEN Board of Directors and the Clinical Guidelines Task Force. Guidelines for the use of parenteral and enteral nutrition in adult and pediatric patients. *JPEN J Parenter Enteral Nutr* 2002; 26: 144. [\[Crossref\]](#)
3. Taylor BE, McClave SA, Martindale R, Warren MM, Johnson DR, Braunschweig C, et al. Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *Crit Care Med* 2016; 44: 390-438. [\[Crossref\]](#)
4. Stroud M, Duncan H, Nightingale J. Guidelines for enteral feeding in adult hospital patients. *Gut* 2003; 52(Suppl VII): vii1-12. [\[Crossref\]](#)
5. Pearce CB, Duncan HD. Enteral feeding. Nasogastric, nasojejunal, percutaneous endoscopic gastrostomy, or jejunostomy: its indications and limitations. *Postgrad Med J* 2002; 78: 198-204. [\[Crossref\]](#)
6. Lochs H, Allison SP, Meier R, Pirlich M, Kondrup J, Schneider S, et al. Introductory to the ESPEN Guidelines on Enteral Nutrition: Terminology, definitions and general topics. *Clin Nutr* 2006; 25: 180-6. [\[Crossref\]](#)
7. Bankhead R, Boullata J, Brantley S, et al. Enteral nutrition practice recommendations. *J Parenter Enteral Nutr* 2009;33(2):122-167. [\[Crossref\]](#)
8. Boullata J, Carrera AL, Harvey L, Escuro AA, Hudson L, Mays A, et al. ASPEN Safe Practices for Enteral Nutrition Therapy. *J Parenter Enteral Nutr* 2017; 41: 15-103. [\[Crossref\]](#)
9. Dietitians Association of Australia. Enteral Nutrition manual for adults in healthcare facilities. 2011.
10. Löser C, Aschl G, Hébuterne X, Mathus-Vliegen EM, Muscaritoli M, Niv Y, et al. ESPEN guidelines on artificial enteral nutrition-percutaneous endoscopic gastrostomy (PEG). *Clin Nutr* 2005; 24: 848-61. [\[Crossref\]](#)
11. American Association of Critical-Care Nurses. AACN Practice Alerts: prevention of aspiration. *Crit Care Nurs* 2016; 36: e20-4. [\[Crossref\]](#)
12. National Collaborating Centre for Acute Care (UK). Nutrition Support for Adults: Oral Nutrition Support, Enteral Tube Feeding and Parenteral Nutrition. London: National Collaborating Centre for Acute Care (UK); 2006 Feb. (NICE Clinical Guidelines, No. 32.) 9, Enteral tube feeding in hospital and the community. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK49253/>
13. Braegger C, Decsi T, Dias JA, Hartman C, Kolacek S, Koletzko B, et al. Practical approach to paediatric enteral nutrition: a comment by the ESPGHAN committee on nutrition. *J Pediatr Gastroenterol Nutr* 2010; 51: 110-22. [\[Crossref\]](#)
14. Winkelmann C, Best K. Formula for Success: Deliver Enteral Nutrition Using Best Practices. *Am Nurs Today* 2009; 4: 18-23.

15. Bourgault AM, Ipe L, Weaver J, Swartz S, O’dea PJ. Development of evidence-based guidelines and critical care nurses’ knowledge of enteral feeding. *Crit Care Nurs* 2007; 27: 17-29. [\[Crossref\]](#)
16. Williams TA, Leslie GD. A review of the nursing care of enteral feeding tubes in critically ill adults: part I. *Intensive Crit Care Nurs* 2004; 20: 330-43. [\[Crossref\]](#)
17. Demirkan K, Bayraktar EA, Enteral Beslenme tüpünden ilaç uygulanmasında ilaç dozaj şekillerinin önemi, *J Turk Soc Intensive Care* 2016; 14: 1-8.
18. Howard P, Jonkers-Schuitema C, Furniss L, Kyle U, Muehlebach S, Ödlund-Olin A, et al. Managing the patient journey through enteral nutritional care. *Clin Nutr* 2006; 25: 187-95. [\[Crossref\]](#)
19. Potter PA, Perry A G, Stockert P, Hall A, *Fundamentals of Nursing*, 9. Edition, Elsevier Health Sciences, Canada, Copyright: 2013.
20. Stroud M, Duncan H, Nightingale J, British Society of Gastroenterology. Guidelines for enteral feeding in adult hospital patients. *Gut* 2003; 52(Suppl 7): vii1-vii12. [\[Crossref\]](#)
21. Williams TA, Leslie GD. A review of the nursing care of enteral feeding tubes in critically ill adults: part II. *Intensive Crit Care Nurs* 2005; 21: 5-15. [\[Crossref\]](#)
22. Resmi gazete, Sağlık Bakanlığı, Hemşirelik Yönetmeliğinde Değişiklik Yapılması Yönetmeliği, 19 Nisan 2011, Sayı: 27910.
23. Best, C. The correct positioning and role of an external fixation device on a PEG. *Nurs Times* 2004; 100: 50-1.
24. Watterson C, Fraser A, Banks M, Isenring E, Miller M, Silvester C, et al. Evidence Based Guidelines for nutritional management of malnutrition in adult patients across the continuum of care. *Nutr Diet* 2009; 66: S1-S34. [\[Crossref\]](#)
25. American Association of Critical-Care Nurses. AACN Practice Alerts: prevention of aspiration. *Crit Care Nurs* 2016; 36: e20-4. [\[Crossref\]](#)
26. Metheny NA. Residual volume measurement should be retained in enteral feeding protocols. *Am J Crit Care* 2008; 17: 62-4. [\[Crossref\]](#)
27. Metheny NA, Schallom L, Oliver DA, Clouse RE. Gastric residual volume and aspiration in critically ill patients receiving gastric feedings. *Am J Crit Care* 2008; 17: 512-9. [\[Crossref\]](#)
28. Roche V. Percutaneous endoscopic gastrostomy. Clinical care of PEG tubes in older adults. *Geriatrics* 2003; 58: 22-6, 28-9.
29. McClave SA, Lukan JK, Stefater JA, Lowen CC, Looney SW, Matheson PJ, et al. Poor validity of residual volumes as a marker for risk of aspiration in critically ill patients. *Crit Care Med* 2005; 33: 324-30. [\[Crossref\]](#)
30. Stroud M, Duncan H, Nightingale J, British Society of Gastroenterology. Guidelines for enteral feeding in adult hospital patients. *Gut* 2003; 52(Suppl 7): vii1-vii12. [\[Crossref\]](#)
31. Dietitians Association of Australia - Nutrition Support Interest Group, Enteral feeding manual for adults in health care facilities. 2011, Dietitians Association of Australia Canberra, Australia. Available from: [http:// daa.asn.au/wp-content/uploads/2011/11/ Enteral-nutrition-manual-Oct-2011.pdf](http://daa.asn.au/wp-content/uploads/2011/11/Enteral-nutrition-manual-Oct-2011.pdf).